

TORADOL (KETOROLAC)

PHARMACOLOGY & MECHANISM OF ACTIONS:

- ❑ Exhibits analgesic, anti-inflammatory, and antipyretic activity.
- ❑ NSAID

INDICATIONS:

- ❑ Management of non-trauma associated pain

CONTRAINDICATIONS:

- ❑ Age > 65 y/o secondary to possible renal insufficiency/failure
- ❑ Pregnancy, active labor, lactating female
- ❑ Patients currently using Coumadin, Plavix, ASA, other NSAIDs or anticoagulants
- ❑ Hypotension
- ❑ History of or active bleeding or ulcer disorder
- ❑ Patients with suspected or confirmed cerebral vascular bleeding
- ❑ Patients in need of immediate surgery
- ❑ Hypersensitivity
- ❑ Nasal polyps
- ❑ Angioedema
- ❑ Asthma

ADMINISTRATION:

	ADULT	PEDIATRIC	
EMT-I	<ul style="list-style-type: none">❑ 15-30 mg IV or 30-60 mg IM❑ Patients ≤50 kgs or 65+ years old, use the lower end range of dose	<ul style="list-style-type: none">❑ Not recommended – contact OLMC	EMT-I

PRECAUTIONS & SIDE EFFECTS:

- ❑ Edema, HTN, N/V, HA GI bleeding
- ❑ Use caution in patients with impaired renal or hepatic function, cardiac decompensation, HTN, COPD, & elderly
- ❑ Not recommended for obstetric analgesia. **CLASS C PREGNANCY CATEGORY**
- ❑ Not recommended for use in children
- ❑ May prolong bleeding time, use caution in patients with coagulation disorders
- ❑ Use caution in orthopedic injuries due to risk of bleeding